



Herbicide

Spoiler® herbicide contains 2,4-D, mecoprop-p and dichlorprop-p.



Three-Way Control of Tough Broadleaf Weeds

Broadleaf weeds don't stand a chance against the combined power of Spoiler® herbicide's three active ingredients. The ultimate three-way combination, Spoiler delivers effective broadleaf weed control with reduced risk to ornamentals and dicamba-sensitive plants.

Apply Spoiler post-emergence to residential and non-residential turfgrasses throughout the spring, summer and fall. It's particularly effective on summer annuals such as oxalis, spurge and knotweed.

KEY PERFORMANCE BENEFITS

- Outstanding post-emergent control of tough broadleaf weeds
- Highest active ingredient loaded three-way available
- Non-dicamba product with equivalent control to conventional three-way mixes
- Low use rates
- No-glug jugs
- Can be mixed with some liquid fertilizers

ACTIVE INGREDIENTS

2,4-D, mecoprop-p
and dichlorprop-p

% ACTIVE INGREDIENTS

33.3%, 8.58% and 8.45%

FORMULATION

liquid

CHEMICAL FAMILY

phenoxy

EPA REGISTRATION

228-513

MODE OF ACTION

auxin growth regulator

RESTRICTED USE

no

SIGNAL WORD

warning

PACKAGING INFORMATION

2 x 2.5 gal
30 gal



LABELED USES

Ornamental turf (*lawns, parks, cemeteries and golf courses*)
Along highways and other non-crop areas
Sod farms

DISEASES CONTROLLED

| | |
|----------------------------|---|
| Alder | Oxalis |
| Artichoke | Parsnip |
| Aster | Pennycress |
| Austrian fieldcress | Pennywort |
| Beggartick | Peppergrass |
| Biden | Pepperweed |
| Bindweed | Pigweed |
| Bitterweed | Plantain |
| Bitter wintercress | Poison hemlock |
| Black medic | Poison ivy |
| Blue lettuce | Poison oak |
| Box elder | Pokeweed |
| Broomweed | Poorjoe |
| Buckhorn | Povertyweed |
| Burdock | Prickly lettuce |
| Bur ragweed | Primrose |
| Buttercup | Puncture vine |
| Carpetweed | Purslane |
| Catnip | Ragweed |
| Chickweed | Red clover |
| Chicory | Red sorrel |
| Clover | Rush |
| Cocklebur | St. Johnswort |
| Coffeebean | Sheep sorrel |
| Coffeeweed | Shepherd's-purse |
| Creeping jenny | Smartweed |
| Croton | Sneezeweed |
| Curly indigo | Southern wild rose |
| Dandelion | Sowthistle |
| Dock | Spanishneedle |
| Dogbane | Spatterdock |
| Dogfennel | Speedwell |
| Elderberry | Spurge |
| English daisy | Spurweed |
| Flea bane (<i>daisy</i>) | Stinkweed |
| Flixweed | Stitchwort |
| Florida pusley | Sumac |
| Frenchweed | Sunflower |
| Galinsoga | Sweet clover |
| Garlic mustard | Tarweed |
| Goatsbeard | Thistle (<i>blessed, bull, Canada, musk, Russian</i>) |
| Goldenrod | Toadflax |
| Ground ivy | Tumbleweed |
| Gumweed | Velvet leaf |
| Hawkweed | Veronica |
| Healall | Vervain |
| Heartleaf drymary | Vetch |
| Hemp | Virginia buttonweed |
| Henbit | Virginia creeper |
| Hoary cress | Wild aster |
| Honeysuckle | Wild carrot |
| Horsetail | Wild garlic |
| Indiana mallow | Wild geranium |
| Ironweed | Wild lettuce |
| Jewelweed | Wild onion |
| Jimsonweed | Wild radish |
| Kochia | Wild rape |
| Knotweed | Wild strawberry |
| Lambsquarters | Wild sweet potato |
| Locoweed | Willow |
| Lupines | Witchweed |
| Mallow | Woodsorrel |
| Marshelder | Wormseed |
| Mexicanweed | Yarrow |
| Morningglory | Yellow rocket |
| Mustard | and other |
| Nettle | broadleaf weeds |
| Orange hawkweed | |



Herbicide

APPLICATION RECOMMENDATIONS

FOR BEST RESULTS

Turf should not be mowed for 1 to 2 days before and after application.

Do not apply if rainfall is expected within 8 hours. Do not irrigate for 24 hours following application.

Reseed no sooner than three to four weeks after application of this product.

APPLICATION USE RATES

Cool-season Grasses (such as bluegrass, fescue and ryegrass):

1.1 to 1.5 fl oz/1,000 sq ft, 3 to 4 pts per acre

Warm-season Grasses (such as bahia, common Bermuda and zoysia):

1.75 to 2.5 pts/maximum in 20 to 230 gallons of water per acre

Bentgrass Turf (putting and bowling greens):

.62 to .92 fl oz/1,000 sq ft, 1.7 to 2.5 pts per acre

Better Choices. Better Business. | www.nufarm.com/us

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.

©2009 Nufarm. Important: Always read and follow label instructions. Spoiler® is a trademark of Nufarm.

09-TO-0082

