



**Australian Pesticides &  
Veterinary Medicines Authority**

**PERMIT TO ALLOW MINOR USE OF AN AGVET CHEMICAL PRODUCT**

**PERMIT NUMBER -PER10459**

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows any person to claim that the product can be used in the manner specified in this permit.

**THIS PERMIT IS IN FORCE FROM 1 JANUARY 2008 TO 31 DECEMBER 2012.**

**Permit Holder:**

**GROWCOM**

**Floor 1**

**385 St Pauls Terrace**

**FORTITUDE VALLEY QLD 4006**

**Persons who can use the product under this permit:**

Persons generally.

## CONDITIONS OF USE

### Product to be used:

BASF BAVISTIN FL SYSTEMIC FUNGICIDE

SPIN FLO SYSTEMIC FUNGICIDE

NUFARM CARBEND FUNGICIDE

FARMOZ HOWZAT SC SYSTEMIC FUNGICIDE

Containing: 500 g/L CARBENDAZIM as their only active constituent.

### Directions for Use:

Crop	Pest	Rate
Custard Apple	<b>Control of fungal diseases caused by</b> <i>Cylindrocladium spp. &amp;</i> <i>Pseudocercospora spp.</i>	Apply at a rate of 50 mL of product per 100 mL of water.

### Safety Directions:

**WARNING:** Contains carbendazin which causes birth defects in laboratory animals. Women of child bearing age should avoid contact with carbendazin.

Harmful if inhaled or swallowed. Will irritate the eyes and skin. DO NOT inhale vapour or spray mist. When opening the container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and a **half-facepiece respirator**. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

### Critical Use Comments:

- Apply a maximum of 4 sprays.
- First spray to be applied at fruit set after flowering. Where previous problems have been experienced apply a second spray 2-4 weeks later.
- If high disease pressure should occur, a further 2 sprays may be applied.
- All sprays must be applied at a minimum interval of 14 days.

**Application:** Apply via a misting machine approximately 2L prepared spray per tree or equivalent 1000L prepared spray per hectare.

**Restraint:** DO NOT apply CARBENDAZIM in combination (tank mixes) with products containing COPPER OXYCHLORIDE.

### Withholding Period:

DO NOT harvest for 3 days after application.

### Jurisdiction:

NSW and QLD only.

**Additional Conditions:**

This Permit provides for the use of a product in a manner other than specified on the approved label of the product. Unless otherwise stated in this permit, the use of the product must be in accordance with instructions on its label.

Persons who wish to prepare for use and/or use the products for the purposes specified in this permit must read, or have read to them, the permit particularly the information included in Details of Permit and Conditions of Permit.

**Export of Produce:**

Exported produce must have appropriate residue tolerance limits established in the importing countries and any residues must not exceed the tolerance limits.

Issued by

Delegated Officer