

ChampION⁺⁺™

FUNGICIDE / BACTERICIDE*



Supercharged Copper for Supercharged Control.

ChampION⁺⁺™ has the smallest, most consistent copper ions (particles) of any WG copper formulation. The result: better coverage and better disease control. And, ChampION⁺⁺ delivers all that with less environmental load than high dose copper products.

SUPERIOR COVERAGE AND CONTROL

- Certified for use in organic crop production
- Small particle size (average = 1.2 micron) results in excellent coverage and disease protection
- Stable WG formula pours easily and disperses quickly
- High density copper product means lower use rates and smaller pack sizes
- Controls plant pathogenetic fungi and bacteria*

*Non public health bacteria



NOW OMRI LISTED®
FOR ORGANIC CROP PRODUCTION

EPA REG. NO.	55146-115
ACTIVE INGR.	copper hydroxide (46.1%)
FORMULATION	water-dispersible granule
CHEM. FAMILY	inorganic
GROUP NO.	M1
MODE OF ACT.	multi-site contact activity
REI	24 hours (greenhouse uses) 48 hours
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	10 x 4 lb, 4 x 10 lb, 1 x 30 lb



Grow a better tomorrow.

ChampION⁺⁺™

FUNGICIDE / BACTERICIDE

KEY USES

Citrus

Field crops; alfalfa, barley, corn, oats, peanut, potato, soybean*, sugarbeet, wheat

Grapes and berries; brambles, blueberry, cranberry, currant gooseberry, kiwifruit, raspberry, strawberry

Hops

Tree crops; almond, apple, apricot, avocado, cherry, filbert, macadamia, nectarine, olive, peach, pear, pecan, pistachio, plum, prune, quince, walnut

Tropical crops; banana, cacao, coffee, guava, litchi, mamey sapote, mango, papaya, passion fruit, plantain, sugar apple

Vegetables; bean, beet, carrot, celery, celeriac, crucifers, cucurbits, eggplant, endive, escarole, garlic, leek, lettuce**, okra, onion, pea, pepper, spinach, tomato, watercress

* Not for use in California

** Not for use in California and Arizona

KEY DISEASES CONTROLLED

(See product label for complete list)

Algal leaf spot
 Alternaria brown spot
 Anthracnose
 Bacterial blast
 Bacterial blight
 Bacterial spot
 Black pitting
 Black rot
 Botrytis blight
 Brown spot
 Cane spot
 Cercospora leaf blight
 Cercospora leaf spot
 Citrus canker (*suppression*)
 Collar rot
 Crown rot
 Downy mildew
 Fire blight
 Gummy stem blight
 Leaf curl
 Leaf rust
 Leaf spot
 Phytophthora brown rot
 Phytophthora foot rot
 Powdery mildew
 Pseudomonas blight
 Scab
 Spanish moss
 Walnut blight
 Yellow rust

USE SUMMARY

RATES: Per Application

Citrus	0.5 lbs/Acre
Field crops	0.5-2.0 lbs/Acre
Grapes and berries	0.75-4.25 lbs/Acre
Hops	0.75-1.5 lbs/Acre
Tree crops	0.5-10.5 lbs/Acre
Tropical crops	0.75-7.75 lbs/Acre
Vegetable crops	0.5-3.5 lbs/Acre

APPLYING SPRAY MIXTURE

The amount of this product applied per acre in concentrate and aerial sprays must be the same as the amount applied per acre in dilute sprays. The required amount of this product must be mixed with enough water to thoroughly cover the crop with spray mixture and is to be applied to the point of runoff for best performance. The volume of water needed depends upon factors such as the spray equipment used, foliage density, size of the crop being sprayed, etc.

CHEMIGATION APPLICATION

Apply this product only through sprinkler irrigation systems including center pivot, motorized lateral move, end tow, traveler, big gun, plastic solid set, or plastic hand move; DO NOT apply this product through any other type of irrigation system. In California, do not apply in systems which contain aluminum parts or components.

See label for complete directions for use.

APPLICATION PRECAUTIONS

Do not apply this product in a spray solution having a pH of less than 6.5 as phytotoxicity may occur.

Environmental conditions such as extended periods of wet weather, acid rain, etc. which alter the pH of the leaf surface may affect the performance of this product resulting in possible phytotoxicity or loss of effectiveness.

Do not tank mix with other pesticides or additives unless the mixture has been determined to be compatible and non-injurious to the crop under your conditions of use.

See label for complete application precautions.



www.nufarm.com/us

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.

©2015 Nufarm. Important: Always read and follow label instructions. ChampION⁺⁺™ is a trademark of Nufarm. OMRI Listed® is a trademark of Organic Materials Review Institute. 15-AG-0009-B