

Platform™



Build Your Rice Weed Control on a Solid Platform™.

Broadleaf weeds can wreak havoc on rice yields and your profits. Platform™ Herbicide from Nufarm offers excellent control of key weeds like alligatorweed, hemp sesbania and morningglory. With its flexible application timing – burndown, pre-flood or post-flood – Platform offers a strong base for your rice weed control program.

KEY PERFORMANCE BENEFITS

- Outstanding control of key broadleaf weeds like alligatorweed, hemp sesbania and morningglory
- Flexible application timing – burndown, pre-flood or post-flood
- Excellent tank-mix partner

EPA REG. NO.	228-520
ACTIVE INGR.	triclopyr (44.4%)
FORMULATION	liquid
CHEM. FAMILY	pyridine
GROUP NO.	4
MODE OF ACT.	synthetic auxin
REI	48 hours
SIGNAL WORD	danger
RESTRICTED	no
PACKAGE SIZE	2 x 2.5 gal



Nufarm

Grow a better tomorrow.

Platform™

KEY USES

Non-irrigation ditch banks
 Aquatic sites such as ponds, lakes, reservoirs, non-irrigation canals, and ditches which have little or no continuous outflow
 Rice

KEY WEEDS CONTROLLED

(See product label for complete list)

RICE

Alligatorweed
 Cocklebur, common
 Dayflower
 Eclipta
 Hemp sesbania
 Jointvetch
 Morningglory
 Redstem
 Rice flatsedge
 Sicklepod
 Texasweed/Mexicanweed
 Water hyssop
 Ricefield bulrush

AQUATIC

Alligatorweed
 American lotus
 American frogbit
 Aquatic sodapple
 Eurasian watermilfoil
 Milfoil species
 Nuphar (*spatterdock*)
 Parrot feather
 Pennywort
 Pickerelweed
 Purple loosestrife
 Waterhyacinth
 Waterlily
 Waterprimrose

USE SUMMARY

RATES

Rice: Use 10.7 to 16 fluid ounces per acre. Best control is achieved with applications to actively growing weeds prior to flowering. Make post-flood applications when weeds are well-emerged above the water surface. Weeds submerged at application will not be controlled.

Floating and emerged aquatic weeds: Apply 0.5 to 2 gallons of this product per acre as a foliar application for control of waterhyacinth, alligatorweed and other susceptible emerged and floating weeds. Apply using surface or aerial equipment. Use higher rates when plants are mature, when the weed mass is dense, or for difficult-to-control species. Retreatment may be necessary to control regrowth and weeds which escaped spray, but do not apply more than 2 gallons of this product per acre per annual growing season.

Submerged weeds: Apply sufficient product per acre (depending on water depth) to obtain a target site concentration of 0.75 to 2 PPM triclopyr acid. Provides control of Eurasian watermilfoil (*Myriophyllum spicatum*) and other susceptible submerged weeds in ponds, lakes, reservoirs, and in non-irrigation canals or ditches that have little or no continuous outflow. Apply using surface or subsurface application. Use higher rates within the range in areas of greater water exchange. Do not apply more than 2.5 PPM acid equivalent of this product per acre per annual growing season.

See label for complete directions for use

